



**UNIVERSITI PUTRA MALAYSIA**

**PREPARATION AND CHARACTERISATION OF CROSS-LINKED  
SAGO STARCH PHOSPHATES**

**FARIDAH BINTI HUSSIN**

**FSMB 1998 18**

**PREPARATION AND CHARACTERISATION OF CROSS-LINKED  
SAGO STARCH PHOSPHATES**

**FARIDAH BINTI HUSSIN**

**MASTER OF SCIENCE  
UNIVERSITI PUTRA MALAYSIA**

**1998**



**PREPARATION AND CHARACTERISATION OF CROSS-LINKED  
SAGO STARCH PHOSPHATES**

**By**

**FARIDAH BINTI HUSSIN**

**Thesis Submitted in Fulfillment of the Requirements for the Degree of  
Master of Science in the Faculty of Food Science and Biotechnology  
Universiti Putra Malaysia**

**September 1998**



## ACKNOWLEDGEMENTS

Alhamdulillah, first of all I would like to express my utmost thanks and gratitude to Almighty Allah S.W.T. who has given me the capability to complete this project and my salawat and salam to His messenger, Prophet Muhammad s.a.w.

I would like to express my most sincere appreciation and deepest gratitude to my supervisor, Dr. Sharifah Kharidah binti Syed Muhammad for her understanding, willingness, generous gift of time and guidance, invaluable advice, constructive suggestions and valuable comments which had helped tremendously in the preparation of this thesis. I am also very grateful for the patience, care and encouragement that she had so generously offered during this study and in reviewing the manuscript.

My heartfelt appreciation and sincere thanks goes to the member of my supervisory committees, Prof. Dr. Yaakob bin Che Man and Prof. Madya Dr. Hasanah binti Mohd Ghazali who had selflessly devoted their time to read and review the manuscript in great detail time after time. Their generous guidance, invaluable comments, advice and suggestions in the preparation of this thesis is deeply appreciated. I would also like to express my sincere thanks for their care and encouragement during the course of my study in UPM and most of all for their warm friendship.

My appreciation and special thanks also goes to my former supervisor, Dr. Mohd. Nasir bin Azudin for his brilliant ideas, technical guidance and advice during this study.



My deepest appreciation and gratitude are accorded specially to En. Dzulkifly Mat Hashim, En. Razali Mustaffa, Pn. Zaulia Othman, En. Amin Ismail, Miss Irene Ch'ng, Dr. Puziah Hashim, Cik Jamaliah Ahmad, En. Azman Abu Yamin, En. Chan Tin Wan, Kak Hus, En. Halim, Dr. Mwanjallah Mwasaru, Miss Malini and all my friends, technicians, laboratory assistants and staff members of the Faculty of Food Science and Biotechnology for their very generous help, care, encouragement and warm friendship which had made my study in UPM a very pleasant and memorable experience.

Most of all, I would like to express my deepest gratitude to my beloved parents, husband, and children for their endless encouragement, love, patience and sacrifices which helped me in my undertakings and to complete this work.

## TABLE OF CONTENTS

	<b>Page</b>
AKNOWLEDGEMENTS .....	ii
LIST OF TABLES .....	vii
LIST OF FIGURES .....	viii
LIST OF PLATES .....	xii
LIST OF ABBREVIATIONS .....	xiii
ABSTRACT .....	xiv
ABSTRAK .....	xvi
<b>CHAPTER</b>	
<b>1      GENERAL INTRODUCTION .....</b>	<b>1</b>
<b>2      LITERATURE REVIEW .....</b>	<b>5</b>
Modified Starches .....	5
Cross-linked Starch .....	8
Manufacturing Process .....	8
Effect of Processing Parameters .....	12
General Reactions .....	15
Types of Starches .....	20
Characteristics and Properties of Cross-linked Starch.....	20
Phosphorus in Starch .....	20
Pasting Profile .....	21
Degree of Substitution (DS).....	26
Sediment Volume .....	27
Paste Clarity .....	28
Water Holding Capacity .....	29
Swelling Power and Solubility .....	31
Scanning Electron Microscopy .....	32
Gel Strength of Starch Paste .....	32
Freeze-thaw Stability .....	33
Starch Damage .....	36
Utilisation .....	36
Sago starch .....	39
Origin .....	39
Extraction of Sago Starch.....	40
Properties and Chemical Compositions .....	40
Utilisation .....	41

### 3      **EFFECT OF VARIOUS PROCESSING PARAMETERS ON THE PHOSPHORUS CONTENT, PASTING PROFILE, AND PASTE CLARITY OF PHOSPHORYLATED SAGO STARCH**

Introduction .....	44
Materials and Methods .....	45
Materials .....	45
Starch Damage .....	46
Total Amylose .....	48
Phosphorylation of Starch .....	49
Phosphorus Content .....	50
Pasting Profiles .....	50
Paste Clarity .....	51
Results and Discussion .....	51
Effect of pH on the Sago Starches Phosphorylated with STMP, STPP, and Mixture of the Two .....	53
Effect of Sodium Tripolyphosphate (STPP) Concentrations .....	63
Effect of Sodium Trimetaphosphate (STMP) Concentrations .....	69
Effect of Sodium Sulfate Concentrations .....	74
Effect of Reaction Time .....	78
Effect of Reaction Temperature .....	83
Effect of Heating Time .....	89
Effect of Heating Temperature .....	92
Conclusions .....	98

### 4      **PHYSICO-CHEMICAL CHARACTERISTICS OF CROSS-LINKED SAGO STARCH COMPARED WITH COMMERCIAL CROSS-LINKED WAXY CORN STARCHES**

Introduction .....	99
Materials and Methods .....	100
Materials .....	100
Preparation of Cross-linked Sago Starch .....	101
Phosphorus Content of Starch .....	101
Degree of Substitution .....	101
Swelling Power and Solubility .....	102
Pasting Profile .....	102
Paste Clarity .....	102
Sediment Volume .....	103
Gel Strength of Starch Paste .....	103
Acid and Heat Stability .....	104
Water Holding Capacity .....	104
Freeze Thaw Stability of Gels .....	105
Scanning Electron Microscopy (SEM) .....	105
Starch Damage .....	106

Results and Discussion .....	106
Phosphorus Content of Starch .....	106
Degree of Substitution .....	108
Swelling Power and Solubility .....	108
Pasting Profiles.....	111
Paste Clarity .....	113
Sediment Volume .....	115
Gel Strength of Starch Paste .....	117
Acid and Heat Stability .....	119
Water Holding Capacity .....	124
Freeze-Thaw Stability of Gels .....	126
Scanning Electron Microscopy (SEM) .....	128
Starch Damage .....	133
Conclusion .....	133
 <b>5      GENERAL CONCLUSION AND RECOMMENDATIONS ..</b>	 136
General Conclusion and Recommendations.....	136
 <b>BIBLIOGRAPHY .....</b>	 139
 <b>BIOGRAPHICAL SKETCH.....</b>	 153



## LIST OF TABLES

Table		Page
1	Disadvantages of unmodified starches .....	6
2	Types of modified starches .....	7
3	Dissociation constant of phosphoric acids .....	16
4	Effect of autoclaving on brookfield viscosity and solubility of starches .....	26
5	Water binding capacities of native and modified starches .....	30
6	Influence of cross-linking on granular swelling power (GSP) of corn and waxy corn .....	32
7	Properties of starches .....	42
8	Physico-chemical composition of sago starch .....	52
9	Phosphorus content of native, cross-linked sago and cross-linked waxy corn starches (Mazaca 3543X and Mazaca 3544X) .	107
10	Degree of substitution of native, cross-linked sago and cross-linked waxy corn starches (Mazaca 3543X and Mazaca 3544X) .	109
11	Water holding capacity of native, cross-linked sago and cross-linked waxy corn starches (Mazaca 3543X and Mazaca 3544X) .	125
12	Starch damage of native sago, cross-linked sago and cross-linked waxy corn starches (Mazaca 3543X and Mazaca 3544X) .	134

## LIST OF FIGURES

Figure	Page
1 Cross-linking to supplement hydrogen bonding in a starch granule .....	9
2 Structure of distarch Glycerol (A) and distarch phosphate .....	9
3 Reactions of sodium tripolyphosphate with starch at alkaline and acidic pH. Slanted arrow indicates the phosphate attacked by a starch hydroxyl .....	17
4 Reactions of sodium trimetaphosphate with starch at alkaline and acidic pH. Slanted arrow indicates the phosphate attacked by a starch hydroxyl .....	18
5 Effect of cross-linking corn starch on brabender viscograph .....	24
6 Effect of cross-linking waxy corn starch on brabender viscograph at pH 5 and 3 .....	23
7 Meaning of DS and MS. On the left, the molecules has only one substituent X, which gives a DS and MS equal to 1. On the right, there are three substituents X, which gives a DS of 1 and a MS of 3.	27
8 Freeze-thaw stability of 5% pastes of various starches .....	35
9 Effect of various initial reaction pH on the total phosphorus content of sago starch phosphates prepared using 2% sodium trimetaphosphate (STMP), 5% sodium tripolyphosphate (STPP), and a mixture of the two .....	54
10 Amylograms of sago starch phosphates prepared by reacting starch at various initial pH levels with 5% sodium tripolyphosphate based on dry starch. All amylograms were obtained at pH 6.5 and 6% starch solids .....	57
11 Amylograms of sago starch phosphates prepared by reacting starch at various initial pH levels with 2% sodium trimetaphosphate based on dry starch. All amylograms were obtained at pH 6.5 and 6% starch solids .....	58
12 Amylograms of sago starch phosphates prepared by reacting starch at various initial pH levels with a mixture of 5% sodium sulfate, 5% sodium tripolyphosphate and 2% sodium trimetaphosphate based on dry starch. All amylograms were obtained at pH 6.5 and 6% starch solids .....	60

13	Paste clarity of sago starch phosphates prepared using 5% sodium tripolyphosphate (STPP), 2% sodium trimetaphosphate (STMP), and a Mixture of the two at various pH levels .....	62
14	Effect of sodium tripolyphosphate concentrations on the total phosphorus content of sago starch phosphates prepared at pH 11...	64
15	Amylograms of sago starch and sago starch phosphates prepared at various concentrations of sodium tripolyphosphate (STPP) based on dry starch at pH 11. All amylograms were obtained at pH 6.5 and 6% starch solids .....	66
16	Effect of sodium tripolyphosphate concentrations on paste clarity of sago starch phosphates prepared at pH 11 based on dry starch...	68
17	Effect of sodium trimetaphosphate concentrations on the total phosphorus content of sago starch phosphates prepared at pH 8...	70
18	Amylograms of sago starch and sago starch phosphates prepared at various concentrations of sodium trimetaphosphate (STMP) based on dry starch at pH 8. All amylograms were obtained at pH 6.5 and 6% starch solids .....	71
19	Effect of sodium trimetaphosphate concentrations on paste clarity of sago starch phosphates prepared at pH 8 based on dry starch ....	73
20	Effect of sodium sulfate concentrations on the total phosphorus content of sago starch phosphates prepared using a Mixture of 2% sodium trimetaphosphate and 5% sodium tripolyphosphate based on dry starch at pH 9.5 .....	75
21	Amylograms of sago starch and sago starch phosphates prepared by reacting starch at various concentrations of sodium sulfate based on dry starch at pH 9.5. All amylograms were obtained at pH 6.5 and 6% starch solids .....	77
22	Effect of sodium sulfate concentrations on paste clarity of sago starch phosphates prepared using 5% sodium tripolyphosphate (STPP) and 2% sodium trimetaphosphate (STMP) at based on dry starch pH 9.5.....	79
23	Effect of various reaction times on the total phosphorus content of sago starch phosphates prepared using a mixture of 2% sodium trimetaphosphate and 5% sodium tripolyphosphate based on dry starch at pH 9.5 .....	80
24	Amylograms of sago starch and sago starch phosphates prepared by reacting starch at various reaction times at pH 9.5. All amylograms were obtained at pH 6.5 and 6% starch solids .....	82

25	Effect of various reaction times on paste clarity of sago starch phosphates prepared using 5% sodium tripolyphosphate (STPP) and 2% sodium trimetaphosphate (STMP) at based on dry starch pH 9.5.....	84
26	Effect of various reaction temperatures on the total phosphorus content of sago starch phosphates prepared using a mixture of 2% sodium trimetaphosphate and 5% sodium tripolyphosphate based on dry starch at pH 9.5 .....	85
27	Amylograms of sago starch and sago starch phosphates prepared by reacting starch at various reaction temperatures at pH 9.5. All amylograms were obtained at pH 6.5 and 6% starch solids .....	87
28	Effect of various reaction temperatures on paste clarity of sago starch phosphates prepared using a mixture of 2% sodium trimetaphosphate and 5% sodium tripolyphosphate based on dry starch at pH 9.5 .....	88
29	Effect of various heating times on the total phosphorus content of sago starch phosphates prepared using a mixture of 2% sodium trimetaphosphate and 5% sodium tripolyphosphate based on dry starch at pH 9.5 .....	90
30	Amylograms of sago starch and sago starch phosphates prepared by reacting starch at various heating times at pH 9.5. All amylograms were obtained at pH 6.5 and 6% starch solids .....	91
31	Effect of various heating times on paste clarity of sago starch phosphates prepared using 5% sodium tripolyphosphate (STPP) and 2% sodium trimetaphosphate (STMP) at based on dry starch pH 9.5.....	93
32	Effect of various heating temperatures on the total phosphorus content of sago starch phosphates prepared using a mixture of 2% sodium trimetaphosphate and 5% sodium tripolyphosphate based on dry starch at pH 9.5 .....	94
33	Amylograms of sago starch and sago starch phosphates prepared by reacting starch at various heating temperatures at pH 9.5. All amylograms were obtained at pH 6.5 and 6% starch solids .....	96
34	Effect of various heating temperatures on paste clarity of sago starch phosphates prepared using a mixture of 2% sodium trimetaphosphate and 5% sodium tripolyphosphate based on dry starch at pH 9.5 .....	97
35	Swelling power and solubility of native sago, sago and waxy corn starches .....	110

36	Amylograms of native sago, cross-linked waxy corn (Mazaca 3543X and Mazaca 3544X) starches.	112
37	Paste clarity of native sago, starches .....	114
38	Sediment volume of native sago, corn starches .....	116
39	Gel strength of native sago, starches .....	118
40	Acid and heat stability of native sago starch. All amylograms were obtained at 6% starch solids .....	120
41	Acid and heat stability of cross-linked sago starch. amylograms were obtained at 6% starch solids .....	121
42	Acid and heat stability of cross-linked waxy corn starch (Mazaca 3543X). All amylograms were obtained at 6% starch solids .....	122
43	Acid and heat stability of cross-linked waxy corn starch (Mazaca 3544X). All amylograms were obtained at 6% starch solids .....	123
44	Freeze-thaw stability of native sago, cross-linked sago and waxy corn starches (Mazaca 3543X and Mazaca 3544X). One cycle is equal to two days of storage .....	127

## LIST OF PLATES

Plate		Page
1	Scanning electron micrographs of native sago starch at 500X and 1,000X magnifications .....	129
2	Scanning electron micrographs of cross-linked sago starch at 450X and 1800X magnifications .....	130
3	Scanning electron micrographs of cross-linked waxy corn starch (Mazaca 3543X) at 550X and 2000X magnifications.....	131
4	Scanning electron micrographs of cross-linked waxy corn starch (Mazaca 3544X) at 850X and 2000X magnifications.....	132

## LIST OF ABBREVIATIONS

AGU	Anhydroglucose unit
BU	Brabender unit
°C	degree Centigrade
DS	Degree of Substitution
dsb	dry starch basis
gm	gram
hr	hour
M	Molar
min	minute (s)
ml	milliliter
nm	nanometer
P	Phosphorus
CFR	Code of Federal Regulation
SIRIM	Standard and Industrial Research Institute of Malaysia
STMP	Sodium trimetaphosphate
STPP	Sodium tripolyphosphate
UV	Ultra Violet
VIS	Visible
μm	micrometer
<	less than
%	percent

Abstract of Thesis Presented to the Senate of Universiti Putra Malaysia in  
Fulfillment of the Requirements for the Degree of Master of Science.

## **PREPARATION AND CHARACTERISATION OF CROSS-LINKED SAGO STARCH PHOSPHATES**

BY

**FARIDAH HUSSIN**

September, 1998

Supervisor : Sharifah Kharidah Bte Syed Muhammad, Ph. D.

Faculty : Food Science and Biotechnology

This project was conducted to study the production of cross-linked sago starch phosphate and to characterise the material produced. Sago starch was phosphorylated at various pHs (from 6 to 11) under a range of phosphate salt concentrations (sodium tripolyphosphate [STPP] [1 to 9%] and sodium trimetaphosphate [STMP] [1 to 4%]) and in the presence of various sodium sulfate concentrations (1 to 9%). The phosphorylation was carried out for 40 to 80 mins at temperatures varying from 27 to 50°C and heating time from 0.5 to 3 hrs at temperatures between 100 to 160°C. It was observed that the reaction pH and concentration of phosphate salts played a significant effect on the phosphorus (P) content, pasting profile, and paste clarity of the sago starch phosphates produced. The phosphorus content was found to increase as the pH and concentration of phosphate salts were increased under all conditions. However, the phosphorus contents were found to be below the standard (STPP [0.4%], STMP [0.04%], and a mixture of STPP and STMP [0.4%]), except when the concentration of STMP was above 1%. The pasting profile showed that at pH 9.5, treatment of sago starch with a



mixture of 5% STPP and 2% STMP yielded the best cross-linked sago starch phosphate where it showed the lowest hot paste viscosity and the highest cold paste viscosity. Paste clarity measurements of the phosphorylated starches indicated that cross-linking had accelerated rapidly with STMP above pH 8, with STPP above pH 9, and with a mixture of the two above pH 6. Paste clarity measurements also showed that cross-linking began to accelerate at 5% and below STPP, and 2% and below STMP when sago starch was phosphorylated at pH 11 and 8, respectively. Judging from the paste properties, phosphorylation of sago starch at 27°C for 1 hr with a mixture of 5% STPP and 2% STMP at pH 9.5 in the presence of 5% sodium sulfate and then heating at 130°C for 2 hr is recommended. Cross-linked sago starch phosphate and commercial cross-linked waxy corn starch (Mazaca 3543X and 3544X) showed similar pasting profile. However, cross-linked sago starch phosphate had higher degree of phosphorylation (phosphorus content), degree of substitution, swelling power and solubility, paste clarity and gel strength. The water holding capacity and sediment volumes were similar to one of the cross-linked waxy corn starches (Mazaca 3544X). Morphology studies showed that these modified starches retained their granule shape with minimal degree of starch damage.

Abstrak tesis yang dikemukakan Kepada Senat Universiti Putra Malaysia  
Sebagai Memenuhi Syarat Keperluan Untuk Ijazah Sarjana Sains

## **PENYEDIAAN DAN PENCIRIAN KANJI SAGU IKATAN-SILANG FOSFAT**

Oleh

**FARIDAH HUSSIN**

September, 1998

Penyelia : Sharifah Kharidah Bte Syed Muhammad, Ph. D.

Fakulti : Sains Makanan dan Bioteknologi

Projek ini dijalankan untuk mengkaji penghasilan dan pencirian kanji sagu fosfat ikatan-silang daripada kanji sagu. Kanji sagu telah difosforilasi pada pH yang berbeza (dari 6 ke 11) dengan menggunakan kepekatan garam-garam fosfat yang berbeza (sodium tripolifosfat [STPP] [1 hingga 9%] dan sodium trimetafosfat [STMP] [1 hingga 4%]) dan dengan kehadiran kepekatan sodium sulfat yang berbeza (1 hingga 9%). Fosforilasi telah dijalankan selama 40 hingga 80 minit pada suhu daripada 27 hingga 50°C dan dipanaskan selama 0.5 hingga 3 jam pada suhu diantara 100 hingga 160°C. Didapati paras pH tindakbalas dan kepekatan garam-garam fosfat menunjukkan kesan yang bermakna keatas kandungan fosforus (P), profil kelikatan, dan kejernihan pes kanji-kanji sagu fosfat yang dihasilkan. Kandungan fosforus didapati bertambah apabila pH dan kepekatan garam-garam fosfat bertambah pada semua keadaan. Walaubagaimanapun, kandungan fosforus didapati di bawah paras piawaian (STPP [0.4%], STMP [0.04%], dan campuran STPP dan STMP [0.4%]), kecuali apabila kepekatan STMP melebihi 1%. Profil kelikatan menunjukkan bahawa kanji sagu yang dirawat dengan campuran 5% STPP dan 2% STMP pada pH 9.5 menghasilkan kanji fosfat yang terbaik dimana ia

menunjukkan kelikatan pes panas yang terendah dan kelikatan pes sejuk yang lebih tertinggi. Pengujian kejernihan pes ke atas kanji yang telah difosforilasi menunjukkan bahawa ikatan-silang telah meningkat dengan cepatnya pada pH lebih daripada 8 dengan STMP, lebih daripada 9 dengan STPP, dan lebih daripada 6 dengan campuran kedua-duanya. Pengukuran kejernihan pes juga menunjukkan bahawa ikatan-silang mula bertambah pada 5% STPP dan ke bawah, dan ke bawah apabila kanji sagu difosforilasikan pada pH 11 dan 8, masing-masing. Merujuk kepada ciri-ciri kelikatan, difosforilasi pada suhu 27°C selama 1 jam dengan menggunakan campuran 5% STPP dan 2% STMP pada pH 9.5 dengan kehadiran 5% sodium sulfat dan kemudiannya dipanaskan pada suhu 130°C selama 2 jam digunakan. Kanji sagu fosfat ikatan-silang dan kanji jagung berlilin ikatan silang (Mazaca 3543X dan 3544X) menunjukkan profil kelikatan yang sama. Walaubagaimanapun, ikatan silang menunjukkan darjah fosforilasi (kandungan fosforus), penggantian, kuasa pengembangan dan kelarutan, kejernihan pes dan kekuatan gel yang lebih tinggi. Kapasiti pegangan air dan isipadu mendapan adalah sama dengan salah satu daripada kanji-kanji jagung berlilin ikatan silang (Mazaca 3544X). Kajian morfologi menunjukkan kanji-kanji yang telah diubahsuai ini tidak berubah bentuk granul asalnya dengan darjah kerosakan yang rendah.

## CHAPTER 1

### GENERAL INTRODUCTION

Cross-linked starches are those products resulting from an intentional reaction of starch with bi- or polyfunctional reagents, which will effect a bond between glucose units on separate chains. They are widely used as thickening agents in sauces and dressing, bakery products, puddings, desserts and custards, soups, and emulsifying agent in many food systems, antiblocking dusting agent, warp sizing of denims and surgical dusting powder in other applications (Swinkels, 1992). The types of cross-linked that are of commercial value are distarch adipates, distarch phosphates, and distarch glycerol. Of these, only distarch phosphates and distarch glycerol are used in making modified food starches (Wurzburg, 1986b). In this study, distarch phosphates will be produced through phosphorylation because it is a method commonly employed for cross-linking starches intended for the food industry and other uses (Koch *et al.*, 1982).

Distarch phosphates are produced when a phosphate radical bridges two starch molecules and in consequence the starch granule is inhibited from swelling and rupture on pasting in water (Radley, 1976). A small amount of cross-linking greatly reduces both the rate and the degree of granule swelling and sensitivity of starch pastes to processing conditions.

Starch can be cross-linked by reaction with various inorganic phosphate salts as well as specially developed organic reagents. The U.S. Food and Drug Administration allows starch for use in foods to be modified with sodium trimetaphosphate (residual phosphorus not exceeding 0.04%), sodium trimetaphosphate and sodium tripolyphosphate (residual phosphorus not exceeding 0.4%) and phosphorus oxychloride (maximum allowable treatment is 0.1% on starch) (CFR, 1991).

Cross-linked starches have been produced mainly in United States, Europe, Japan and Australia. These products are mostly derived from corn and tapioca starch. As a result, technical information regarding the manufacturing process and the various processing parameters are based on modifications of these starches (Wurzburg, 1986; Rutenberg and Solarek, 1984; Kerr and Cleveland, 1959)

In Malaysia, modified starches are imported for utilization in the local food industries and other applications. Referring to the statistics on import and re-export of merchandises prepared by the Malaysian Industrial Development Authority (MIDA), from January 1995 to August 1996, this country has imported 31,286,116.63 tonnes of modified starches which is equivalent to RM64,911,550.00. United States, Italy, Thailand, Australia, France and Germany supply these modified starches. To date, there is only one local factory producing commercial cross-linked starch either for local utilization or for export. This factory is producing cross-linked starch using waxy corn starch which is imported from Australia. Therefore, if cross-linked starch can be manufactured using one of the local indigenous starches, this

will not only save revenue but also diversify the usage of such starch. One such possible indigenous starch which Malaysia is the largest producer and exporter of is sago starch (Azudin and Lim, 1991).

Sago palm occurs throughout Malaysia and it is substantially common in fresh water swamp areas along some river courses and on peat soils (Othman, 1991). Sago starch which is extracted from sago palm has been produced primarily in Sarawak. The main areas of sago starch production in Sarawak are Mukah, Dalat and Igan of the Sibu division. During the last decade, Sarawak has been exporting about 25,000 - 30,000 tonnes of sago starch annually to Peninsular Malaysia, Singapore, Japan and other parts of the world. This monopoly is basically due to the well established sago cultivation areas available and the large scale production of sago starch practiced in this state (Zulpilip *et al.*, 1991). In addition, over the past five years there has been a tremendous improvement in starch extraction and processing technology producing high quality sago starch. Azudin *et al.* (1989) in their evaluation of the local starch have shown that sago starch produced by the modern sago factories are well above the quality specifications of the Standard and Industrial Research Institute of Malaysia (SIRIM) standards for sago starch.

Sago starch has been utilized mainly in the production of 'bee hoon' (or vermicelli), manufacture of monosodium glutamate, glucose, maltose, dextrose, and fish and prawn crackers (Zulpilip *et al.*, 1991). There has been no report or literature on the suitability of sago starch for the production of modified starch especially cross-linked starch. Therefore, the objectives of this study were:

1. To establish the optimum conditions for phosphorylating sago starch
2. To determine the physico-chemical characteristics of the cross-linked sago starch, and to compare them with that of commercial cross-linked waxy corn starches

## **CHAPTER 2**

### **LITERATURE REVIEW**

#### **Modified Starches**

Modified starch is a native starch treated in such a way as to modify one or more of its original physical or chemical properties (Swinkels, 1992). Modification is carried out because unmodified starches cannot be tailored-made and very often are not the best product in a particular application or process as swell with ease and rupture with minimum abuse to produce weak-bodied, cohesive pastes or undesirable gels (Wurzburg, 1986a). The main disadvantages of unmodified starches are shown in Table I.

Modifications of native starches are carried out in order to overcome these disadvantages and thus expand the usefulness of starch for a great number of industrial applications (Wurzburg, 1986a). They are designed to change and improve one or more of the properties as appropriate for a specific application (IFI, 1993) (Table 2).



**Table 1: Disadvantages of unmodified**

1. **Inconsistent viscosity.** Due to the effects of climatic condition of the plant.
2. **Susceptibility to acid attack.** Many food are acidic. A food product made using an unmodified starch will be unstable. Unmodified starch is degraded by the acid hydrolysis.
3. **Susceptibility to shear.** An unmodified starch will lose its viscosity due to the mechanical action of stirring. Therefore, the viscosity of the final product depending upon how long it has been stirred.
4. **Set-back and syneresis.** In solution the hydrogen bonding increases with time as thermal motion brings various chains into close enough proximity for further hydrogen bonding. As a colloidal sol 'sets back' into a gel on standing, the process continues with occluded water molecules. The gel to leave a rubbery solid surrounded by the expressed liquid. This is accelerated if the product is frozen then thawed.
5. **Solubility.** A problem arises in any process requiring high solids content due to the limit on the amount that can be 'dispersed' as a colloid, imposed by the macromolecular nature of starch.

---

Source : Rothwell and Garner (1986)